

a pharmaceutically acceptable acid addition salt or a stereochemically isomeric form thereof, wherein

C1 cont
R¹ and R² are each independently selected from hydrogen; hydroxy; amino; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; Ar¹; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C₁₋₆alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxyC₁₋₆alkyloxy, carboxyl, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxycarbonyl and thienyl; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₆alkyl)aminoC₁₋₄alkylidene;

B1
R³ is hydrogen, Ar¹, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxycarbonyl; and

R⁴, R⁵, R⁷ and R⁸ are each independently selected from hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethoxy;

R⁶ is aminocarbonyl; or

--- ???
L is C₁₋₁₀ alkyl; C₃₋₁₀alkyl; C₃₋₁₀alkyl; C₃₋₇cycloalkyl; or C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C₁₋₆alkylcarbonyl, phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C₁₋₆alkylcarbonyl; and, Ar¹ is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl.